



Fenton, I., Pearson, P.N., Dunkley Jones, T., Farnsworth, A. J., Lunt, D. J., Markwick, P., & Purvis, A. (2016). The impact of Cenozoic cooling on assemblage diversity in planktonic foraminifera. *Philosophical Transactions B: Biological Sciences*, 371, [20150224]. <https://doi.org/10.1098/rstb.2015.0224>

Peer reviewed version

Link to published version (if available):
[10.1098/rstb.2015.0224](https://doi.org/10.1098/rstb.2015.0224)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

The impact of Cenozoic cooling on assemblage diversity in planktonic foraminifera

Isabel S. Fenton^{1,2*}, Paul N. Pearson³, Tom Dunkley Jones⁴, Alexander Farnsworth⁵, Daniel J. Lunt⁵, Paul Markwick⁶ and Andy Purvis^{1,2}

¹Department of Life Sciences, Natural History Museum, Cromwell Road, London, U.K.

²Department of Life Sciences, Imperial College London, Silwood Park campus, Ascot, U.K.

³School of Earth and Ocean Sciences, Cardiff University, Cardiff, U.K.

⁴School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham U.K.

⁵School of Geographical Sciences & Cabot Institute, University of Bristol, Bristol, U.K.

⁶Getech Group plc., Elmete Hall, Elmete Lane, Leeds, U.K.

*Corresponding author: isabel.fenton@nhm.ac.uk

Key words: Planktonic foraminifera, Eocene, Latitudinal diversity gradient, global climate model

Abstract

The Cenozoic planktonic foraminifera (calcareous zooplankton) have arguably the most detailed fossil record of any group. The quality of this record allows models of environmental controls on macroecology, developed for Recent assemblages, to be tested on intervals with profoundly different climatic conditions. These analyses shed light on the role of long-term global cooling in establishing the modern latitudinal diversity gradient – one of the most powerful generalizations in biogeography and macroecology. Here we test the transferability of environment-diversity models developed for modern planktonic foraminiferal assemblages to the Eocene epoch (~56 to 34 Mya), a time of pronounced global warmth. Environmental variables from global climate models are combined with Recent environment-diversity models to predict Eocene richness gradients, which are then compared with observed patterns. The results indicate the modern latitudinal diversity gradient – lower richness towards the poles – developed through the Eocene. Three possible causes are suggested for the mismatch between statistical model predictions and data in the early Eocene: the environmental estimates are inaccurate, the statistical model misses a relevant variable, or the intercorrelations among facets of diversity – e.g. richness, evenness, functional diversity – have changed over geological time. By the late Eocene environment-diversity relationships were much more similar to those found today.

Key words: Planktonic foraminifera, Eocene, Latitudinal diversity gradient, global climate model

Introduction

Ecologists and palaeontologists have long been interested in the causes of diversity patterns found in many taxonomic groups, but the two communities often approach such questions in different ways [1, 2]. Ecologists are able to obtain diversity estimates across the globe, but usually lack a temporal perspective [e.g. 3], whereas palaeontologists typically have a more limited spatial resolution at any given point in time [4]. Each approach has its strengths and limitations. If evolutionary processes such as speciation, immigration and extinction have been important for structuring Recent diversity patterns [e.g. 5, 6], then incorporating fossil data into studies of these patterns is likely to be important [7]. Although phylogenies of extant species contain some information about the rates of these processes, they lack direct information about ancestors or extinct species limiting the inferences that can be drawn [8, 9]. Combining ecological and palaeontological approaches is likely to aid our understanding of diversity patterns [1, 2, 10, 11].

The latitudinal diversity gradient (LDG) is one of the most widely studied of all diversity patterns. It occurs in a very broad range of habitats and taxa [12, 13]. Despite being well-described, there is little consensus on whether the LDG has one dominant underlying cause or reflects multiple causes acting in concert [12, 14, 15]. Community species richness is often found to be correlated with environmental variables [e.g. 15, 16], but these correlations do not prove the true causes have been identified [17]. Instead these correlations could have a separate, underlying cause. Using an independent dataset can improve support for the hypothesis that the important variables have indeed been captured.

There are three ways to obtain such an independent dataset. The first is to test the predictions in a different region. This is not possible for global diversity studies where the whole clade is included in the original analysis. The second method transfers the model to a different clade [e.g. 18, 19, 20]. There are, however, limits to how well any model would be expected to transfer among clades, given that all clades have some degree of difference in ecological traits and tolerances. It is thus hard to separate a mismatch caused by ecological differences from that caused by an imperfect model. The third method transfers the model to a time period with substantially different environmental conditions, a concept often used in global climate modelling [e.g. 21]. Typically macroecologists lack a deep-time temporal perspective, limiting such transfers to a few decades [e.g. 22, 23]. Such short timescales are unlikely to be informative for global diversity patterns; global environmental conditions will not vary sufficiently and they are likely to be too short for macroevolutionary processes such as speciation to act.

Most study systems, even those with excellent fossil records, are so spatiotemporally and / or taxonomically incomplete that advanced methods are needed to estimate even simple diversity measures such as species richness [24] and biogeographic history [25]. Without the use of such methods, spatiotemporal diversity patterns derived from the fossil record of most extant clades are too fragmentary for a robust test of hypotheses developed from the modern biota. Two approaches have been used to obtain sufficient species-level data for diversity gradient studies: (i) the study of a small number of sites with very high quality data are compared to a known diversity gradient, e.g. that in the Recent [e.g. 26]; or (ii) all known fossils from a time period are grouped into latitudinal bins [e.g. 27].

1 The planktonic foraminifera (PF) provide a rare study system where neither approximations nor
2 complex methods are necessary: assemblage diversity estimates can be obtained consistently
3 across large spatial and temporal scales. As such their importance for deepening our
4 understanding of macroecological patterns is now being recognised [2, 16]. These calcareous
5 open-ocean protists have the most complete fossil record of any clade [28, 29], with species
6 having at least an 81% chance of being recorded from any given 1Myr (million year) time bin
7 [30], equivalent to the best genus-level completeness for macroinvertebrates [31]. Deposition in
8 deep marine environments, where planktonic foraminifera are typically recorded, is more
9 continuous than either shallow marine or terrestrial environments [32], reducing the need to
10 untangle the correlation between rock record and diversity that occurs in the other settings [e.g.
11 33, 34]. Assemblages from the deep-ocean are thus more equivalent across spatial and
12 temporal scales than terrestrial or shallow marine assemblages [35].

13 In this paper we start by summarising the geographic patterns of Recent planktonic foraminiferal
14 diversity and the macroecological models used to determine the drivers of that diversity. We
15 then make use of the clade's exceptional fossil record, interrogating a new collation of Eocene
16 assemblage data, to determine diversity patterns in the Eocene. We combine our present day
17 statistical models with palaeoenvironmental data from climate models to derive estimates of
18 Eocene diversity patterns which are then compared to observed Eocene diversity to determine
19 the transferability of Recent macroecological models.

20 Recent planktonic foraminiferal diversity

21 Foraminifera are unicellular zooplankton with a test or 'shell', usually made of calcium
22 carbonate. This test consists of a series of chambers which are added progressively as the
23 organism grows; holes, or foramina, allow movement of cytoplasm between these chambers
24 [36]. Foraminifera show two main lifestyles: benthic foraminifera live on the seafloor, while
25 planktonic foraminifera (PFs) float passively in the ocean. PFs can be further split into
26 macroperforates or microperforates, depending on the size of the pores in the test. Here we
27 focus on macroperforate species which are less susceptible to dissolution than the
28 microperforates, are more frequently identified in palaeontological studies, comprise a single
29 and dominant clade, and have a better understood taxonomy and ecology [29].

30 The tests of fossil PFs include all the taxonomically-diagnostic morphological characters used to
31 classify living specimens. This makes Recent and fossil PF morphospecies fully comparable,
32 contrasting with the situation in most other taxa where non-comparability of taxonomic
33 categories hampers comparisons of fossil and Recent diversity [37]. Molecular analyses (using
34 SSU rRNA) however suggest some extant morphospecies should be split into cryptic or semi-
35 cryptic genetic species, with slightly different ecological and biogeographical preferences [e.g.
36 28, 38, 39]. To date, around half of modern morphospecies have been sequenced, with the
37 presence of cryptic diversity varying across the clade [39]. For example, genetic studies of
38 *Orbulina universa* identified three pseudo-cryptic species inhabiting different hydrological
39 conditions [40], whereas the *Trilobatus sacculifer* complex contains a wide range of
40 morphologies while showing no evidence for genetic variability [41, 42]. As possible cryptic
41 species within a morphospecies are mostly geographically separated, occupying slightly
42 different ecological or environmental niches [e.g. 39], they are likely to have more impact on

1 global diversity estimates than on species richness counts at a given location. It is also
2 important to note the cryptic species problem is not unique to PFs, and the ratio of possible
3 genetic species to morphospecies is in line with that seen in other taxa [e.g. 43]. Although SSU
4 rRNA underestimates true diversity in many other groups [44], it evolves much more rapidly in
5 PFs [45].

6 PFs are well sampled both in the Recent and through the Cenozoic, as a result of their use as
7 stratigraphic marker fossils and for palaeoenvironmental proxies [46]. Core-top marine sediment
8 samples are routinely analysed for oceanographic studies, including collecting census data for
9 calcareous microplankton. Sample preparation and data collection are usually broadly
10 consistent between studies, in marked contrast to much terrestrial biodiversity monitoring.
11 Typically, the sample is sieved at 150µm and 300 individuals are counted [following 47]. The
12 MARGO dataset, used to provide the Recent diversity data in this analysis, contains
13 approximately 4000 assemblage counts of Recent core-top samples [48; Supplementary Figure
14 1]. The data include relative abundances of all observed species; only morphospecies that are
15 genuinely absent, very rare or very small (and so passing through the sieve) will not be
16 recorded in such an assemblage count.

17 These data can be visualised as species distribution maps (e.g. Figure 1). Based on these
18 distributions, extant PF diversity is split into six faunal provinces each with a distinct community
19 [36, 49]. Few species are found exclusively in one faunal province; instead such provinces are
20 defined by the dominant species. Provinces tend to correspond to the major ocean gyre
21 systems, running roughly parallel to lines of latitude. High-productivity upwelling regions,
22 however, are distinct from other sites at the same latitude. In the modern oceans, these regions
23 tend to contain relatively higher abundances of *Neogloboquadrina*, *Globigerina bulloides* and
24 some genetic variants of *Globigerinella siphonifera* [39].

25 Assemblage count data have long been used to calculate a wide range of site-level diversity
26 measures based on taxonomic diversity [50]. Data compilations of species' traits and
27 evolutionary relationships [29] now also allow functional and phylogenetic diversity to be
28 calculated. Recent developments in automating trait measurements allow studies in the spatial
29 patterns of morphological variance [51]. As these core top samples represent death
30 assemblages, they are not directly comparable to a single census of live specimens.
31 Communities living at different depths or in different seasons are averaged in the sediment.
32 Lateral transport of individuals before deposition may also lead to a certain amount of spatial
33 averaging [52], but this is thought to be relatively minimal for PFs [28].

34 Here we focus on four diversity measures to characterise different aspects of assemblage
35 diversity: rarefied species richness, Simpson's evenness, mean evolutionary age and functional
36 richness. Rarefied species richness is a sample-size-independent estimate of site-level species
37 richness [53]; samples were rarefied to 275 individuals. Evenness quantifies whether
38 assemblages are dominated by a small number of species. To remove the correlation with
39 species richness which can occur in evenness measures (e.g. Shannon diversity, Simpson's
40 index), we calculate Simpson's evenness calculated as Simpson's index divided by species
41 richness; low values imply a few species dominate [54]. Functional richness is defined as the

amount of functional space filled by the community and measured using Villéger *et al.*'s [55] FRic metric. The functional traits included in this measure are the test structure and size, as well as life habits of the species (presence/absence of spines and symbionts, ecogroup and morphogroup [29], depth habitat, dissolution susceptibility). The mean evolutionary age (MEA) is the (abundance-weighted) average age of the species lineages present in an assemblage [56]. To calculate this measures accurately requires a fossil record of the quality of the PFs. Molecular phylogenies will always assign sister species the same age, defining age as the split between two extant species [57]. Morphospecies ages in the fossil record can also be biased: PF morphospecies are used as zone fossils, encouraging the fine-scale division of species by time [46], even in the absence of cladogenetic events, meaning that the appearance of a morphospecies may reflect a pseudospeciation [e.g. 58, 59]. We therefore use evolutionary species (lineages), as determined by Aze *et al.* [29], for calculating the ages. These originate at cladogenesis and terminate with extinction of the lineage, but can persist through speciation if the speciation has a budding (offshoot) rather than bifurcation (equal splitting) pattern [60-62]. A younger MEA implies that species in an assemblage have arisen more recently. However, a community's MEA does not imply that it has existed for that length of time, or that any speciation occurred *in situ*.

The latitudinal variation of these four diversity measures for the Recent PFs were explored in detail by Fenton *et al.* [56]. Whereas the peak in terrestrial species richness typically occurs at the equator [12], PFs – in common with many other marine taxa [13, 63] – have two peaks in diversity at about $\pm 20^\circ$ latitude. These peaks correspond to the oligotrophic subtropical gyres, whereas the equator is subject to upwelling of deep, nutrient rich water, creating more eutrophic conditions. Functional diversity, like richness, is low at high latitudes; but it approaches its maximum more rapidly, showing little variation below 40° . This levelling out implies functional redundancy in tropical regions. Temperate and high latitude sites typically have more equal abundances than in tropical sites, i.e. have a higher Simpson's evenness. Upwelling regions are less speciose, with less even assemblages. Although MEA shows relatively little change with latitude the oldest assemblages are in subpolar waters while polar assemblages are the youngest (but are very species-poor).

Drivers of Recent diversity

Explanations of diversity patterns can be split into ecological, evolutionary, historical or statistical causes [e.g. 64, 65]. Ecological causes assume that the LDG results from processes acting over relatively short spatial and temporal scales, on factors such as suitable habitat, available energy, competition between species and dispersal limitations. For example, Briggs [66] suggests richness is higher at warmer temperatures as there is sufficient energy for specialisation on relatively low-energy food sources, which at cooler temperatures would not suffice. Evolutionary explanations for the LDG invoke different rates of speciation, extinction or immigration at different latitudes [e.g. 5, 66]. The 'out of the tropics' model, for example, assumes that taxa preferentially originate in the tropics, subsequently expanding into higher latitudes. The metabolic theory of ecology relates high tropical richness to faster speciation at higher temperatures [67]. Historical explanations implicate a region's history through geological time – either changes in the physical properties, such as plate tectonics driving climate change, or the occurrence of contingent events, such as meteorite impacts. The Mid-Domain Effect

(MDE), a statistical explanation, states that if species' ranges are placed randomly into an area with hard boundaries, a gradient will develop with a central peak. The width of this peak depends on range sizes, being wider when ranges are small. As the poles act as hard boundaries, the MDE will produce an equatorial peak in diversity, i.e., an LDG [68]. Alternative versions of the MDE suggest that it could act on an environmental gradient such as temperature [20].

Macroecological models of foraminiferal diversity have mostly focused on species richness, often in the Atlantic Ocean. Rutherford *et al.* [16] reported nearly 90% of species richness variation in the Atlantic can be explained by sea surface temperature (SST); other variables, such as temperature at depth, productivity and salinity, did not significantly improve their models (Table 1). They suggested SST influences diversity by controlling the vertical partitioning of the water column, with the associated creation of distinct niches, but did not test this hypothesis. A similar study [13] of multiple taxa and all the oceans found multiple factors were independently significant predictors of richness (Table 1), although temperature was still the most important. The correlation between species richness and temperature has been found to hold for the last three million years [69]

It is becoming increasingly recognised that incorporating multiple facets of diversity (e.g. taxonomic, phylogenetic, functional) improves understanding of the origins of diversity patterns [70, 71]. Other studies have related more aspects of community structure to a set of environmental variables (Table 1). Morey *et al.* [72] used canonical correspondence analysis to identify environmental correlates of community structure but did not interpret their results in terms of drivers of diversity. Fenton *et al.* [56] tested many of the ecological and evolutionary hypotheses proposed to explain PF diversity by relating the four diversity measures above to a set of environmental variables chosen to capture ocean temperature, structure, productivity and seasonality (see Table 1; Supplementary Figure 2). Their models find support for a combination of ecological and evolutionary models of diversity. Although sea-surface temperature explains the largest portion of diversity in all four diversity measures, observed relationships do not match metabolic theory of ecology or mid-domain models predictions [56]. Historical models are thought to be less significant for PFs due to the absence of dispersal limitation in this clade [73]. These results suggest the diversity patterns of PFs cannot be explained by any one environmental variable or proposed mechanism but reflect multiple processes acting in concert.

Descent into the Icehouse

Temporal comparisons are most powerful in testing macroecological models when the predictive time differs markedly in environmental conditions from that of the initial analysis [23]. Previous work suggests that the richness-temperature relationship in PFs has remained constant for 3Myr, though the diversity patterns have changed alongside climate [69]. Here we consider more aspects of diversity and environment, and attempt to transfer the relationship over a much longer time period. The Eocene (56 to 34 Mya; million years ago) had a broadly similar palaeogeographic configuration of the major ocean basins as the Recent, and a phylogenetically and ecomorphologically comparable PF community, but a very different global climate, so provides a strong test of model transferability. By considering time-slices through the Eocene it is possible to interrogate the effects of the cooling trend from early Eocene conditions

of Cenozoic peak warmth through to late Eocene climates, immediately prior to the onset of the present “icehouse” climate state [74]. The mid to late Eocene saw changes in the distribution of ocean fronts and regions of productivity [75, 76], as well as in the ocean nutricline and the structuring of planktonic niches [77, 78]. By the late Eocene, global cooling had produced biogeographically distinct high-latitude communities [79].

Observed diversity through the Eocene

Eocene PFs occupied similar niches to extant species, being globally distributed with ecologies dependent on depth habitat, hydrography and mode of life [80]. Many Eocene taxa have analogues in the Recent, such as the digitate morphology in *Clavigerinella jarvisi* (Eocene) and *Beella digitata* (extant); isotopic analyses indicate a deep-dwelling habit for both species [81]. Some Eocene groups and traits however are no longer represented in the biota. Most species with the ‘muricate’ wall structure, characteristic of the acarininids, went extinct in the Eocene, and this morphology was finally lost in the Oligocene [29]. The *Hantkenina* lineage initially occupied a unique warm deep-water niche [78], which was lost when it migrated permanently into a shallower habitat [82]. Previous studies of Eocene diversity have mainly focussed on the global picture [e.g. 30, 83], although Boersma and Premoli Silva [79] analysed site-level data from the Atlantic to suggest the onset of an LDG in species richness by the end of the Eocene.

The Eocene PF assemblage data compiled for this project were obtained from a range of sources, and have never previously been collated in one place (Supplementary information). The basis of the data was the NEPTUNE dataset, a repository of records from the ocean drilling programmes [84, 85]. This dataset is now 15 years out of date, so it was supplemented by an extensive literature search including more recent drilling programme publications. The taxonomy of Eocene PFs, including full synonymy lists, was revised by Pearson *et al.* [80], and subsequently combined into a look-up table to ensure taxonomic consistency [as used in 29]. Where possible the most recent calibrations of planktonic foraminiferal zones [46] were used to date the samples. Palaeolocations of sites were calculated using the Getech PIC plate model [following 86] which determines locations at the time at which samples were deposited. This plate reconstruction is consistent with that used for the environmental data from the climate models described in the next section. Sites that showed high levels of dissolution were excluded to prevent systematic bias [28, 87], as were sites where only a fraction of the species were identified, as occurs where the primary purpose was biostratigraphy. Where there were multiple estimates of diversity for the same site within a time interval, the mean diversity observed at a site was chosen to represent the total diversity. (Similar results are obtained if the maximum value is used; not shown). The complete dataset contains 78 sites with a reasonable spatial coverage (Supplementary Figures 3 & 4). To assess latitudinal gradients, the Eocene was divided into three time intervals: early (56-47.8 Myr), middle (47.8-38 Myr) and late (38-33.9 Myr).

We calculated the same four diversity metrics for each site as in the Recent, except that in-sample species richness was used in place of rarefaction-based richness. Both in the Recent and the Eocene, some studies report data as percentage abundance rather than counts of individual. However the Eocene dataset is too small to include only studies with count data. Where it could be calculated, rarefied richness was strongly correlated with, though usually

1 lower than, observed in-sample richness (linear model: $r^2 = 0.69$, $n = 158$, $p < 10^{-15}$,
2 Supplementary Figure 5).

3 Generalized additive models (GAMs) [88] of assemblage diversity with latitude in each sub-
4 epoch show that the LDG developed during the Eocene (Figure 2, Supplementary Figure 6).
5 The significance of these diversity changes does not depend on any individual site
6 (Supplementary Table 1). The gradient changed shape significantly through this time period.
7 Similar to today, middle and late Eocene diversity peaked at 20°-30° latitude. Equatorial species
8 richness may have been lower than in the present-day, although equatorial sites in the early
9 Eocene are sparse. The Simpson's evenness gradient did not change significantly through the
10 Eocene, although it was consistently higher than in the present-day, implying less dominance by
11 a few species. Functional richness did not change significantly through the Eocene (Figure 2).
12 High-latitude functional richness dropped slightly from the early to the middle Eocene, indicating
13 loss of some functional groups, whether through extinction or range shift. Despite the drop in
14 species richness at lower latitudes in the mid-late Eocene there is no associated drop in the
15 functional richness. This disconnect implies that the LDG is not solely driven by niche
16 availability.

17 Mean evolutionary age (MEA) changed significantly through the Eocene, developing a latitudinal
18 gradient. This change was driven by an ageing of high-latitude assemblages, suggesting new
19 species were less able to enter, persist in or dominate these communities. Low-latitude
20 assemblages had similar average ages through time, despite the ageing of the clade as a
21 whole, implying the loss of older species and/or the gain of new ones. The early Cenozoic
22 rebound from the Cretaceous/Paleogene mass extinction sets an upper limit on MEA that
23 increases over time. Periods with higher speciation rates, such as the Palaeocene/Eocene
24 boundary and the mid-Eocene [30], reduce global average MEA estimates.

25 Predicted diversity in the Eocene

26 Testing the transferability of the Recent macroecological models of PF diversity requires
27 estimates of Eocene palaeoenvironment. These are combined with the Recent diversity-
28 environment models to generate predicted diversity values for comparison with the Eocene
29 assemblage data. In this transfer we focus only on the richness-environment model, as richness
30 has the strongest relationship with environment in the Recent [pseudo- $r^2 = 0.89$, see 56].
31 Eocene environmental estimates are inevitably far less certain than modern observational data
32 in both accuracy and spatial coverage. Proxy-based estimates, e.g. of sea surface temperature
33 [e.g. 89, 90], have sparse spatial coverage and so are insufficient for spatially-explicit global
34 models, although we qualitatively compare trends inferred from these estimates with observed
35 richness data. For spatially-explicit global estimates – maps of predicted diversity – we instead
36 use output from an Atmosphere-Ocean General Circulation Model (AOGCM), which provides
37 global coverage of a range of variables. Specifically, we use the HadCM3L model, as used in as
38 used in Lunt *et al.* [91] and Inglis *et al.* [86]. This model has a spatial resolution of 2.5° by 3.75°
39 (latitude and longitude, respectively), with 19 atmospheric and 20 oceanic levels. For further
40 information on these models see Inglis *et al.* [86] and Lunt *et al.* [92]. The model has been
41 evaluated against proxy data [86, 89].

1 We considered two suites of simulations from the AOGCM which represent two potential long-
2 term drivers for Eocene climatic cooling: (i) global CO₂ decline and (ii) plate tectonic changes to
3 oceanic gateways. The first suite – the CO₂ suite – held palaeogeography constant (in an early
4 Eocene configuration), altering only the CO₂ concentrations [early Eocene – 1680 ppmv, middle
5 Eocene – 1120 ppmv, late Eocene – 560 ppmv: 91]. The second suite – the tectonic suite – kept
6 CO₂ levels at 1120ppmv throughout while changing the land-mass configurations [early Eocene
7 – Ypresian; middle Eocene – Bartonian; late Eocene – Priabonian: 86]. These simulations allow
8 us to investigate the individual impact of CO₂ and tectonic change on species richness.
9 Unfortunately model simulations which combine both CO₂ and tectonic change through the
10 Eocene are currently unavailable.

11 The predicted species richness for each suite of simulations was calculated by combining the
12 palaeoenvironmental data with an adapted version of the richness-environment model
13 developed for the Recent [56]. The list of variables included in this adapted statistical model can
14 be found in Table 1; they are the same as the Recent except that productivity and oxygen stress
15 are not predicted in the AOGCM, or otherwise available for the Eocene, so are omitted. The
16 oceanic water mass is also excluded as it is not comparable between the two time periods.
17 GAMs were used rather than spatial autoregressive models, as the former are less likely to give
18 extreme values when extrapolating beyond the range of the data used to fit the model. To
19 mitigate spatial autocorrelation in the GAMs, a two-dimensional smooth of latitude and longitude
20 was included [88]. All statistical modelling analyses were performed in R version 3.0.3 [93].

21 Richness was estimated for each grid cell of the environmental data in each time period. The
22 goodness-of-fit between observations and the model predictions for the corresponding grid cells
23 was quantified using the root mean squared error (RMSE), the average absolute departure of
24 points from the fitted values. For each simulation's predictions, the AIC of how much the
25 observed differed from the predicted diversity was calculated [94]. We compared the goodness-
26 of-fit of the simulations in the CO₂ and the tectonic suites, using Δ AIC to determine whether the
27 richness data provides stronger support for either simulation. Models with Δ AIC > 4 are taken as
28 having 'considerably less' support than the minimum-AIC model [95]. As SST is the most
29 important single correlate of species richness in the Recent [16; Supplementary Figure 2],
30 temperature estimates are likely to have the greatest influence on the richness estimates. To
31 test whether temperature by itself is sufficient to explain the observed richness, a similar GAM
32 containing only temperature was estimated from the Recent data and used to estimate Eocene
33 richness. The RMSE values for this statistical model were also calculated for comparison with
34 the full model.

35 Figure 3 compares the predicted and the observed Eocene species richness; Table 2 gives the
36 corresponding RMSE and Δ AIC values. These results suggest that the tectonic and CO₂
37 simulations differ little in their predictions for the latitudinal patterns of richness. The fit at high
38 latitudes is best by the end of the Eocene; at low latitudes, the middle Eocene produces a
39 slightly better fit (Table 2). However the overall shape is correctly predicted for the middle and
40 late Eocene, but not for the early Eocene. The tectonic simulations fit significantly better in the
41 Early Eocene, but the CO₂ simulations provide the markedly better fit by the end of the Eocene.
42 Re-running the analysis using only the predicted relationship between temperature and richness

1 produces only marginally better fit throughout the Eocene (Table 2), with the early Eocene
2 shape still incorrect (Figure 3).

3 **Possible reasons for the mismatch between predicted and observed** 4 **diversity**

5 The lack of fit between assemblage data and model predictions suggests that the relationship
6 between foraminiferal richness and environmental drivers observed today transfers imperfectly
7 to the Eocene, especially the early Eocene (Figure 3, Table 2). There are three possible
8 reasons for this mismatch. First, the AOGCM estimates of the palaeoenvironmental variables
9 used in this analysis could be incorrect. If so, even a statistical model that correctly captured the
10 environment-richness relationship would appear to fail in the Eocene. Second, the variables
11 included in the model for the Recent could have failed to capture the true drivers of diversity, so
12 any attempt to transfer into a very different past would automatically fail. Third, the relationship
13 between environment and richness could have fundamentally changed since the Eocene. In this
14 case, however correct the statistical model is in the Recent, it would not predict the past
15 accurately. We consider each of these possibilities in turn.

16 **Estimates of Eocene environment**

17 In the early Eocene, the assemblage data suggests the LDG is basically flat from the equator to
18 $\pm 60^\circ$, whereas the model predicts a strong gradient in richness. This prediction arises from the
19 strong latitudinal gradient in temperature present even in the early Eocene in both the
20 simulations (Supplementary Figure 7). Proxy temperature records, by contrast, indicate a
21 weaker [89, 90] or minimal [86] early Eocene temperature gradient, which is inferred to have
22 strengthened through the Eocene [86, 96, 97]. The proxy-based climate estimates are
23 qualitatively more compatible with the observed richness than are the (spatially explicit) climate
24 models we have used. Recent efforts to resolve this proxy-climate model mismatch, based
25 around the choice of parameters associated with clouds and aerosols are promising [98-100],
26 and have the potential to resolve the offsets in observed and predicted richness patterns.
27 However, it is outside the scope of this study to test this possibility.

28 **Explanatory variables for the global diversity model**

29 Several environmental variables – oxygen stress, productivity and ocean – had to be excluded
30 from our analysis, despite having been shown to be significant predictors of Recent diversity
31 [56; Supplementary Figure 2]. Levels of oxygen stress in the Eocene are not clear, although
32 there may have been a more pronounced oxygen minimum in the upper water column due to
33 faster microbial respiration of sinking organic matter [77, 101]. There is some evidence that
34 productivity was high during parts of the Eocene [102, 103]. The differences between Eocene
35 and Recent in the tectonic plate position and the locations of ocean gateways mean that ocean
36 effects on diversity are likely to have altered. Again we cannot test this significance of these
37 missing variables, as they are not currently available from the global climate models.

38 Many environmental variables are strongly correlated with latitude, and consequently, with each
39 other [indeed, this is essentially why the driving mechanisms for the LDG remains a topic of
40 debate: 15]. This correlation makes identifying the true explanatory variables difficult. For
41 example, temperature is known to be important, but in the present day there are strong

1 correlations between mean annual, mean summer and maximum temperature at a site. Each
2 could be used equally well in Recent statistical models, although any one of them might be the
3 variable actually driving richness. If the correlations between these variables have changed,
4 then choosing the wrong variable could lead to a statistical model with a poorer fit in the
5 Eocene. It is very challenging to identify these 'true' variables without a deeper understanding of
6 the ecological response of the foraminifera to their environment.

7 **Diversity-environment relationships**

8 The third explanation suggests the response variable, rather than the explanatory variables,
9 does not fully capture the underlying diversity. Although species richness is often used as a
10 shorthand for biodiversity [15, 104], no single number can adequately capture all facets of
11 biodiver[105]. Diversity measures are often expected to be intercorrelated [106], but the
12 correlations can vary among study systems. For example, although functional diversity usually
13 rises with species richness, there are exceptions [107, 108]. Species richness is often the
14 simplest metric to measure, but may not be the most informative for understanding community
15 structure. The mechanisms that structure communities act upon the ecological similarities,
16 differences and interactions between species, not on numbers of species [109], and the
17 relationships between species richness and other diversity metrics could change through time or
18 space as well as differing among taxonomic groups or ecological guilds. If the relationship
19 between species richness and 'true' diversity (i.e., the idealised measure of diversity that is
20 determined by the environmental drivers) changes, a model relating richness to environment will
21 not transfer, even if the underlying relationships between 'true' diversity and environment are
22 unchanged.

23 To explore this possibility, we undertook a multivariate analysis of six diversity measures, to
24 characterise their intercorrelations in the Recent and the combined Eocene data set. These
25 diversity measures are the four already described (species richness, Simpson's evenness,
26 mean evolutionary age and functional richness) and two closely related measures (mean
27 morphological age and Simpson's diversity). The mean morphological age is the (abundance-
28 weighted) average age of the morphospecies present in an assemblage. As several of the
29 variables were non-normal, we used a robust principal components analysis (rPCA), which
30 scales the data using the median and the median absolute deviation, not the mean and
31 standard deviation [110, 111]. To test whether these two rPCAs differ significantly, we compare
32 the observed PCA similarity [112] to a null distribution obtained of PCA similarity scores
33 obtained from 1000 randomization trials in which the assemblages were randomly divided into
34 two groups having the observed sample sizes.

35 The first three axes of the Recent rPCA explain 94% of the variance; for the Eocene the first
36 three explain 83% of the variance. The rPCA results for the Recent and the Eocene (Figure 4;
37 Supplementary Figure 8) show the relationship between different aspects of diversity have
38 indeed changed through time. For example species richness is strongly correlated with
39 functional richness in the Recent but not in the Eocene. The differences are more pronounced
40 for rPCA2 and rPCA3 than for rPCA1. Assemblages dominated by the same ecogroup are
41 strongly clustered in the Recent, but not in the Eocene. The randomisation test suggests that
42 the Eocene and Recent rPCA structures are near-significantly more different than would be

1 expected by chance (PCAsim = 0.87, significance = 0.071). If we further split the data into each
2 individual time period, there is a suggestion that the diversity structure becomes more similar to
3 the Recent through the Eocene (Supplementary Table 2). However the Eocene datasets are too
4 small to reliably assess the significance of these relationships. A consequence of this apparent
5 change in diversity structure is that, if the drivers of diversity in fact control some facet of
6 diversity other than species richness, then observed species richness could be only a
7 secondary response: the most commonly-used measure of biodiversity may be an
8 epiphenomenon, albeit an extremely useful one for making comparisons within the domain
9 where the biodiversity dimensionality is broadly constant.

10 **Conclusions**

11 Planktonic foraminifera are unusual in allowing the study of how assemblage diversity changes
12 over large spatial and temporal scales. Our analyses identify the 'Descent into the Icehouse' –
13 the global cooling trend through the Eocene – as a key period for the development of the
14 latitudinal diversity gradient within the clade. This analysis suggests the richness-environment
15 relationships seen in the Recent first appeared in the mid-late Eocene, even though the modern
16 PF faunas and provinces did not develop until the Miocene [30, 113]. Our results highlight the
17 Eocene itself, rather than the Eocene-Oligocene boundary, as the time during which the current
18 relationships between environmental drivers and diversity – at least in this clade – became
19 established. An alternative way of viewing the similarity through time in the richness-
20 environment relationship is that PF diversity may respond relatively quickly to environmental
21 change (although this conclusion is tentative pending a comprehensive analysis of assemblage
22 diversity through the entire Cenozoic). Their lack of dispersal limitation [e.g. 39] could be key to
23 this apparently rapid response. In groups with more limited dispersal, the spatial and temporal
24 patterns of speciation and extinction [e.g. 5] may be more important in structuring diversity
25 gradients.

26 Richness-environment relationships appear to have been different in the early Eocene, but we
27 cannot distinguish among several possible explanations: poor estimates of environmental
28 variables (suggested by the mismatch between proxies and GCMs), missing environmental
29 variables, or a fundamental change in the structure of Eocene diversity (shown by the rPCA
30 results and suggested by the extinctions that occur through this time period). However, the
31 similar pattern of high-latitude mismatches when comparing modelled richness with observed
32 richness, and climate model output with climate proxy data suggest that improved models of
33 high-latitude greenhouse climates may be critical to resolving these patterns of biodiversity and
34 climate.

35 **Author contributions:** ISF and AP conceived the project. ISF collated the Eocene PF data and
36 ran the analysis, DJL, PM and AF provided the palaeoenvironment data and the palaeo-
37 coordinates. ISF, PNP, TDJ and AP contributed to the drafting of the manuscript. All authors
38 commented on the manuscript and gave final approval for publication.

39 **Acknowledgements:** We would like to thank Gordon Inglis and Gavin Foster for promoting
40 useful discussion of the palaeoenvironmental data. Also Tom Ezard, Moriaki Yasuhara and an
41 anonymous reviewer for providing helpful comments on the manuscript.

- 1 **Data accessibility:** Eocene diversity data: NHM databank, (<http://dx.doi.org/10.5519/0036305>).
2 For more information see Supplementary information. Recent diversity data available from
3 MARGO: <http://doi.pangaea.de/10.1594/PANGAEA.841194>
4 **Competing interests:** We have no competing interests.
5 **Funding statement:** This work was supported by the NERC funded “Descent into the
6 Icehouse” project (NERC grant NE/I005595/1) and NERC grants NE/M003736/1 and
7 NE/K014757/1.

1 Figure 1. Percentage abundance of *Menardella menardii* in the MARGO; black points indicate
2 absence.

3 Figure 2. Latitudinal trends in diversity of macroperforate planktonic foraminifera through the
4 Eocene; the Recent is added for comparison. GAM smooths are used to highlight the general
5 trends for each time period.

6 Figure 3. Observed and predicted Eocene diversity. Top panels show predictions from the full
7 GAM. Bottom panels show predictions from a temperature only model. Black points / lines are
8 the observed data. Purple (tectonic) and orange (CO₂) points are the model predictions for
9 individual grid cells, with smooths to show the latitudinal trend (continuous lines: northern
10 hemisphere; dashed lines: southern hemisphere).

11 Figure 4. Recent and Eocene rPCA relationships, coloured by the dominant ecogroup [29] at a
12 site. Eco1, open ocean mixed-layer tropical/ subtropical, with symbionts; eco2, open ocean
13 mixed-layer tropical/subtropical, without symbionts; eco3, open ocean thermocline; eco4, open
14 ocean sub-thermocline; eco5, high-latitude. Diversity measures: SR, species richness; FRic,
15 functional richness; MEA, mean evolutionary age; MMA, mean morphospecies age.

16

1

2

Category	Variable	Effect	Study
Energy input	Mean annual SST	Mid-temperature peak	1, 3, 4, 6, 7
	Annual solar irradiance	Positive	1
	MDE on SST	Significant	2, 5
Vertical temperature structure	Mixed-layer depth	Mid-depth peak	1, 6, 7
	Mixed-layer depth variation	None	1
	10°C depth	Mid-depth peak	6, 7
	Temperature at 150 m	Mid-temperature peak	1
Seasonal assemblages	SST variation	None	1, 3, 4, 6, 7
	Salinity variation	Negative	6, 7
Productivity	Mean log Productivity	Mid-productivity peak / none	3, 4, 6
	Mean annual chlorophyll-a	Significant	1, 3
	1% light depth	None	1
	Dissolved nitrate	Significant	1, 3
	Phosphate	Significant	3
Stress	Mean Salinity	Mid-salinity peak	1, 3, 6, 7
	Oxygen stress	Negative	1, 4, 6
Ocean currents	Mean annual topography	Significant	1
	Mean geostrophic current velocity	None	1
	SST slope	Positive	4
Geography	Ocean	None / Significant	4, 6
	Coastline length	Negative	4

	Water depth	Significant	3
Ecological	Temperature niche breadth	Significant	5
Evolution	Geographical origin	Significant	2
Other	Dissolution (when sites with significant dissolution are removed)	None	3, 6, 7
	Density	Significant	3


1 Table 1. Variables used in studies of global diversity in Recent planktonic foraminifera. 1,
2 Rutherford et al (1999), used polynomial regressions; 2, Brayard et al (2005), used a bioclimatic
3 model; 3, Morey et al (2005), used a Canonical Correspondence Analysis; 4, Tittensor et al
4 (2010), used spatial autoregressive models; 5, Beaugrand et al (2013), used a bioclimatic
5 model; 6, Fenton et al (in review), used spatial autoregressive models; 7, this study.

6


Diversity measure	Model	Early Eocene	Middle Eocene	Late Eocene
Mean species richness - Full	Tectonic	7.04 (0)	5.93 (0)	6.27 (+ 7.69)
	CO ₂	7.65 (+ 28.0)	6.72 (+ 2.60)	5.41 (0)
Mean species richness - Temperature	Tectonic	5.76 (0)	5.39 (0)	5.25 (+ 9.91)
	CO ₂	6.18 (+ 23.3)	6.16 (+ 0.21)	5.01 (0)

7 Table 2. Root mean squared error, RMSE, with Δ AIC reported in brackets, of the different
8 models in this analysis.

9




1
2 Figure 1. Percentage abundance of *Menardella menardii* in the MARGO; black points indicate
3 absence.



1 Figure 2. Latitudinal trends in diversity of macroperforate planktonic foraminifera through the
 2 Eocene; the Recent is added for comparison. GAM smooths are used to highlight the general
 3 trends for each time period.

4



5

6 Figure 3. Observed and predicted Eocene diversity. Top panels show predictions from the full
 7 GAM. Bottom panels show predictions from a temperature only model. Black points / lines are
 8 the observed data. Purple (tectonic) and orange (CO₂) points are the model predictions for
 9 individual grid cells, with smooths to show the latitudinal trend (continuous lines: northern
 10 hemisphere; dashed lines: southern hemisphere).




Figure 4. Recent and Eocene rPCA relationships, coloured by the dominant ecogroup [29] at a site. Eco1, open ocean mixed-layer tropical/ subtropical, with symbionts; eco2, open ocean mixed-layer tropical/subtropical, without symbionts; eco3, open ocean thermocline; eco4, open ocean sub-thermocline; eco5, high-latitude. Diversity measures: SR, species richness; FRic, functional richness; MEA, mean evolutionary age; MMA, mean morphological age.

1 **Supplementary Tables**

	Species richness	Species richness 2	Simpson's evenness	Mean Evolutionary Age	Functional richness
Full model	0.0317	0.0164	0.9127	0.00445	0.432
Min.	0.0081	0.0043	0.5407	0.0000313	0.2104
Median	0.0437	0.0171	0.9127	0.00445	0.4319
Mean	0.0416	0.0170	0.8981	0.00498	0.4366
Max.	0.0719	0.0237	0.9965	0.0272	0.6344


2 Supplementary Table 1. P-values from testing whether the diversity-latitude relationship differs
3 significantly between the three Eocene time periods. These values are calculated by comparing
4 a GAM of diversity against latitude where each time interval (i.e. early Eocene, middle Eocene,
5 late Eocene) has a different intercept and slope, with a GAM where the intercepts differ but the
6 slopes are fixed. For all the diversity columns except the second all time-intervals are treated
7 separately. The second species richness column gives the results of combining the middle and
8 late Eocene slopes. For the rows, the full model shows the results with the entire dataset. The
9 remaining rows relate to jackknifing, i.e. testing the reliance of the results on any one point.
10 They show the significance of this comparison when each dataset is excluded in turn. P-values
11 <0.05 imply that the more complex model (i.e. separate slopes and intercepts) is necessary to
12 capture the complexity of the data.

13


	Early	Middle	Late	Recent
Early	x	0.831 (0.212)	0.838 (0.265)	0.816 (0.075)
Middle	x	x	0.935 (0.938)	0.832 (0.227)
Late	x	x	x	0.856 (0.370)

14 Supplementary Table 2. PCA similarity scores for each of the paired comparisons of the periods.
15 The numbers in brackets are the significance of these differences.

Supplementary Figures



Supplementary Figure 1. Distribution of sites for the extant diversity. Data from MARGO [48].




1 Supplementary Figure 2. A comparison of the relative explanatory power of the groups of explanatory
2 variables for the models of the four response variables. Error bars show 1sd and represent the variation
3 associated with removing the replication within each 1 degree square. Figure from [56].




4
5 Supplementary Figure 3. Distribution of sites for the Eocene diversity. Data collated from the literature
6 for this study.

7


Early Eocene




Middle Eocene




Late Eocene




Supplementary Figure 4. The palaeolocations of sites for each period (cf. Supplementary Figure 3 which shows the present day locations). Present day outlines are added for ease of interpretation. Recent has 2389 sites; Early Eocene has 32 sites; Middle Eocene has 26 sites; Late Eocene has 23 sites.




- 1
- 2 Supplementary Figure 5. Comparison of rarefied (black) and normal species richness for those Eocene
- 3 sites with abundance counts. Rarefied richness was calculated for 275 individuals. The lines join the
- 4 species richness and rarefied richness for sites.
- 5



26 Supplementary Figure 6. The GAM smooths and the data points for the diversity measures both
 27 in the Eocene and the Recent (see Figure 2). The dashed lines represent the standard errors.



Supplementary Figure 7. The sea-surface temperature gradient with latitude for the Eocene and the Recent, for the two different climate models tested.



Supplementary Figure 8. Recent and Eocene rPCA relationships, axes 1 and 3. The colours represent the dominant ecogroup [29] at a site: eco1, open ocean mixed-layer tropical/ subtropical, with symbionts; eco2, open ocean mixed-layer tropical/ subtropical, without symbionts; eco3, open ocean thermocline; eco4, open ocean sub-thermocline; eco5, high-latitude.

- [1] Fritz, S.A., Schnitzler, J., Eronen, J.T., Hof, C., Böhning-Gaese, K. & Graham, C.H. 2013 Diversity in time and space: Wanted dead and alive. *Trends Ecol. Evol.* **28**, 509-516.
- [2] Yasuhara, M., Tittensor, D.P., Hillebrand, H. & Worm, B. 2015 Combining marine macroecology and palaeoecology in understanding biodiversity: microfossils as a model. *Biol Rev*, n/a-n/a. (doi:10.1111/brv.12223).
- [3] Orme, C.D.L., Davies, R.G., Burgess, M., Eigenbrod, F., Pickup, N., Olson, V.A., Webster, A.J., Ding, T.-S., Rasmussen, P.C., Ridgely, R.S., et al. 2005 Global hotspots of species richness are not congruent with endemism or threat. *Nature* **436**, 1016-1019.
- [4] Brewer, S., Jackson, S.T. & Williams, J.W. 2012 Paleoecoinformatics: applying geohistorical data to ecological questions. *Trends in Ecology & Evolution* **27**, 104-112. (doi:<http://dx.doi.org/10.1016/j.tree.2011.09.009>).
- [5] Jablonski, D., Roy, K. & Valentine, J.W. 2006 Out of the tropics: Evolutionary dynamics of the latitudinal diversity gradient. *Science* **314**, 102-106. (doi:10.1126/science.1130880).
- [6] Jablonski, D., Belanger, C.L., Berke, S.K., Huang, S., Krug, A.Z., Roy, K., Tomasovych, A. & Valentine, J.W. 2013 Out of the tropics, but how? Fossils, bridge species, and thermal ranges in the dynamics of the marine latitudinal diversity gradient. *Proc. Natn. Acad. Sci.* **110**, 10487–10494. (doi:10.1073/pnas.1308997110).
- [7] Mannion, P.D., Upchurch, P., Benson, R.B.J. & Goswami, A. 2014 The latitudinal biodiversity gradient through deep time. *Trends Ecol. Evol.* **29**, 42-50.
- [8] Purvis, A. 2008 Phylogenetic Approaches to the Study of Extinction. *Annu Rev Ecol Evol S* **39**, 301-319. (doi:10.2307/30245165).
- [9] Quental, T.B. & Marshall, C.R. 2010 Diversity dynamics: molecular phylogenies need the fossil record. *Trends Ecol. Evol.* **25**, 434-441.
- [10] Seddon, A.W.R., Mackay, A.W., Baker, A.G., Birks, H.J.B., Breman, E., Buck, C.E., Ellis, E.C., Froyd, C.A., Gill, J.L., Gillson, L., et al. 2014 Looking forward through the past: Identification of 50 priority research questions in palaeoecology. *J Ecol* **102**, 256-267. (doi:10.1111/1365-2745.12195).
- [11] Benton, M.J. 2015 Exploring macroevolution using modern and fossil data. *Proc. R. Soc. Lond. B* **282**.
- [12] Hillebrand, H. 2004 On the generality of the latitudinal diversity gradient. *Am. Nat.* **163**, 192-211.
- [13] Tittensor, D., Mora, C., Jetz, W., Lotze, H., Ricard, D., Berghe, E. & Worm, B. 2010 Global patterns and predictors of marine biodiversity across taxa. *Nature* **466**, 1098-1101. (doi:10.1038/nature09329).
- [14] Lawton, J.H. 1999 Are there general laws in ecology? *OIKOS* **84**, 177.
- [15] Gaston, K.J. 2000 Global patterns in biodiversity. *Nature* **405**, 220-227.
- [16] Rutherford, S., D'Hondt, S. & Prell, W. 1999 Environmental controls on the geographic distribution of zooplankton diversity. *Nature* **400**, 749-753.
- [17] Rangel, Thiago Fernando L.V.B., Diniz-Filho, José Alexandre F. & Colwell, Robert K. 2007 Species richness and evolutionary niche dynamics: a spatial pattern-oriented simulation experiment. *Am. Nat.* **170**, 602-616. (doi:10.1086/521315).
- [18] Barraclough, T.G., Vogler, A.P. & Harvey, P.H. 1998 Revealing the factors that promote speciation. *Phil. Trans. R. Soc. B* **353**, 241-249. (doi:10.1098/rstb.1998.0206).
- [19] Mittelbach, G.G., Steiner, C.F., Scheiner, S.M., Gross, K.L., Reynolds, H.L., Waide, R.B., Willig, M.R., Dodson, S.I. & Gough, L. 2001 What is the observed relationship between species richness and productivity? *Ecology* **82**, 2381-2396.
- [20] Beaugrand, G., Rombouts, I. & Kirby, R.R. 2013 Towards an understanding of the pattern of biodiversity in the oceans. *Global Ecol. Biogeog.* **22**, 440-449. (doi:10.1111/geb.12009).
- [21] Tindall, J., Flecker, R., Valdes, P., Schmidt, D.N., Markwick, P. & Harris, J. 2010 Modelling the oxygen isotope distribution of ancient seawater using a coupled ocean-atmosphere GCM: Implications for reconstructing early Eocene climate. *Earth Planet. Sci. Lett.* **292**, 265-273.

- [22] Kharouba, H.M., Algar, A.C. & Kerr, J.T. 2009 Historically calibrated predictions of butterfly species' range shift using global change as a pseudo-experiment. *Ecology* **90**, 2213-2222. (doi:10.1890/08-1304.1).
- [23] Rapacciuolo, G., Roy, D.B., Gillings, S., Fox, R., Walker, K. & Purvis, A. 2012 Climatic Associations of British Species Distributions Show Good Transferability in Time but Low Predictive Accuracy for Range Change. *PLoS ONE* **7**, e40212. (doi:10.1371/journal.pone.0040212).
- [24] Starrfelt, J. & Liow, L.H. this volume How many dinosaurs were there? True richness estimated using a Poisson sampling model (TRiPS). *Philosophical Transactions of the Royal Society B: Biological Sciences*.
- [25] Silvestro, D., Bacon, C.D. & Antonelli, A. this volume Effects of climate change on global angiosperm diversification inferred from the fossil record. *philosophical Transactions of the Royal Society B: Biological Sciences*.
- [26] Archibald, S.B., Bossert, W.H., Greenwood, D.R. & Farrell, B.D. 2010 Seasonality, the latitudinal gradient of diversity, and Eocene insects. *Paleobiology* **36**, 374-398. (doi:10.1666/09021.1).
- [27] Mannion, P.D., Benson, R.B.J., Upchurch, P., Butler, R.J., Carrano, M.T. & Barrett, P.M. 2012 A temperate palaeodiversity peak in Mesozoic dinosaurs and evidence for Late Cretaceous geographical partitioning. *Global Ecol. Biogeog.* **21**, 898-908. (doi:10.1111/j.1466-8238.2011.00735.x).
- [28] Kucera, M. 2007 Planktonic foraminifera as tracers of past oceanic environments. In *Developments in Marine Geology* (eds. C. Hillaire-Marcel & A. de Vernal), pp. 213-262, Elsevier.
- [29] Aze, T., Ezard, T.H.G., Purvis, A., Coxall, H.K., Stewart, D.R.M., Wade, B.S. & Pearson, P.N. 2011 A phylogeny of Cenozoic macroperforate planktonic foraminifera from fossil data. *Biol. Rev.* **86**, 900-927. (doi:10.1111/j.1469-185X.2011.00178.x).
- [30] Ezard, T.H.G., Aze, T., Pearson, P.N. & Purvis, A. 2011 Interplay between changing climate and species' ecology drives macroevolutionary dynamics. *Science* **332**, 349-351. (doi:10.1126/science.1203060).
- [31] Foote, M. & Sepkoski, J.J. 1999 Absolute measures of the completeness of the fossil record. *Nature* **398**, 415-417.
- [32] Lloyd, G.T., Pearson, P.N., Young, J.R. & Smith, A.B. 2012 Sampling bias and the fossil record of planktonic foraminifera on land and in the deep sea. *Paleobiology* **38**, 569-584. (doi:10.1666/11041.1).
- [33] Smith, A.B., Lloyd, G.T. & McGowan, A.J. 2012 Phanerozoic marine diversity: Rock record modelling provides an independent test of large-scale trends. *Proc. R. Soc. Lond. B.* (doi:10.1098/rspb.2012.1793).
- [34] Benton, M.J. & Emerson, B.C. 2007 How did life become so diverse? The dynamics of diversification according to the fossil record and molecular phylogenetics. *Palaeontology* **50**, 23-40. (doi:10.1111/j.1475-4983.2006.00612.x).
- [35] Behrensmeyer, A.K., Kidwell, S.M. & Gastaldo, R.A. 2000 Taphonomy and paleobiology. *Paleobiology* **26**, 103-147. (doi:10.1666/0094-8373(2000)26[103:TAP]2.0.CO;2).
- [36] Hemleben, C., Spindler, M. & Anderson, O.R. 1989 *Modern Planktonic Foraminifera*, Springer-Verlag; 363 p.
- [37] Forey, P.L., Fortey, R.A., Kenrick, P. & Smith, A.B. 2004 Taxonomy and fossils: A critical appraisal. *Phil. Trans. R. Soc. B* **359**, 639-653. (doi:10.1098/rstb.2003.1453).
- [38] Huber, B.T., Bijma, J. & Darling, K. 1997 Cryptic Speciation in the Living Planktonic Foraminifer *Globigerinella siphonifera* (d'Orbigny). *Paleobiology* **23**, 33-62. (doi:10.2307/2401156).

- [39] Darling, K.F. & Wade, C.M. 2008 The genetic diversity of planktic foraminifera and the global distribution of ribosomal RNA genotypes. *Marine Micropaleontology* **67**, 216-238. (doi:DOI 10.1016/j.marmicro.2008.01.009).
- [40] de Vargas, C., Norris, R., Zaninetti, L., Gibb, S.W. & Pawlowski, J. 1999 Molecular evidence of cryptic speciation in planktonic foraminifers and their relation to oceanic provinces. *Proc. Natn. Acad. Sci.* **96**, 2864-2868. (doi:10.1073/pnas.96.6.2864).
- [41] André, A., Weiner, A., Quillévéré, F., Aurahs, R., Morard, R., Douady, C.J., de Garidel-Thoron, T., Escarguel, G., de Vargas, C. & Kucera, M. 2013 The cryptic and the apparent reversed: Lack of genetic differentiation within the morphologically diverse plexus of the planktonic foraminifer *Globigerinoides sacculifer*. *Paleobiology* **39**, 21-39. (doi:10.1666/0094-8373-39.1.21).
- [42] Spezzaferri, S., Kucera, M., Pearson, P.N., Wade, B.S., Rappo, S., Poole, C.R., Morard, R. & Stalder, C. 2015 Fossil and genetic evidence for the polyphyletic nature of the planktonic foraminifera "Globigerinoides", and description of the new genus *Trilobatus*. In *PloS one* (p. e0128108).
- [43] Agapow, P.M., Bininda-Emonds, O.R.P., Crandall, K.A., Gittleman, J.L., Mace, G.M., Marshall, J.C. & Purvis, A. 2004 The impact of species concept on biodiversity studies. *Q. Rev. Biol.* **79**, 161-179. (doi:10.1086/383542).
- [44] Tang, C.Q., Leasi, F., Obertegger, U., Kieneke, A., Barraclough, T.G. & Fontaneto, D. 2012 The widely used small subunit 18S rDNA molecule greatly underestimates true diversity in biodiversity surveys of the meiofauna. *Proc. Natn. Acad. Sci.* **109**, 16208-16212. (doi:10.1073/pnas.1209160109).
- [45] Kucera, M. & Darling, K.F. 2002 Cryptic species of planktonic foraminifera: their effect on palaeoceanographic reconstructions. *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences* **360**, 695-718. (doi:10.1098/rsta.2001.0962).
- [46] Wade, B.S., Pearson, P.N., Berggren, W.A. & Pälike, H. 2011 Review and revision of Cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical time scale. *Earth Sci. Rev.* **104**, 111-142.
- [47] CLIMAP. 1976 The Surface of the Ice-Age Earth. *Science* **191**, 1131-1137. (doi:10.1126/science.191.4232.1131).
- [48] Kucera, M., Rosell-Mele, A., Schneider, R., Waelbroeck, C. & Weinelt, M. 2005 Multiproxy approach for the reconstruction of the glacial ocean surface (MARGO). *Quaternary Sci. Rev.* **24**, 813-819.
- [49] Bé, A.W.H. & Tolderlund, D.S. 1971 Distribution and ecology of planktonic foraminifera. In *The Micropaleontology of Oceans* (eds. B.M. Funnell & W.R. Riedel), pp. 105-150, Cambridge University Press.
- [50] Berger, W.H. & Parker, F.L. 1970 Diversity of Planktonic Foraminifera in Deep-Sea Sediments. *Science* **168**, 1345-1347. (doi:10.1126/science.168.3937.1345).
- [51] Hsiang, A., Elder, L. & Hull, P.M. this volume Towards a morphological metric of assemblage dynamics in the fossil record: a test case using planktonic foraminifera. *Philosophical Transactions of the Royal Society B: Biological Sciences*.
- [52] Bé, A.W.H. & Hutson, W.H. 1977 Ecology of planktonic foraminifera and biogeographic patterns of life and fossil assemblages in the Indian Ocean. *Micropaleontology* **23**, 369-414.
- [53] Hurlbert, S.H. 1971 The nonconcept of species diversity: A critique and alternative parameters. *Ecology* **52**, 577-586. (doi:10.2307/1934145).
- [54] Simpson, E.H. 1949 Measurement of diversity. *Nature* **163**, 688.
- [55] Villéger, S., Mason, N.W.H. & Mouillot, D. 2008 New multidimensional functional diversity indices for a multifaceted framework in functional ecology. *Ecology* **89**, 2290-2301.
- [56] Fenton, I.S., Pearson, P.N., Dunkley Jones, T. & Purvis, A. in review Environmental predictors of site-level coretop diversity in Recent planktonic foraminifera. *PLoS ONE*.

- [57] Pigot, A.L., Owens, I.P.F. & Orme, C.D.L. 2012 Speciation and extinction drive the appearance of directional range size evolution in phylogenies and the fossil record. *PLoS Biol* **10**, e1001260. (doi:10.1371/journal.pbio.1001260).
- [58] Stanley, S.M. 1979 *Macroevolution: Pattern and Process*. San Francisco, Freeman.
- [59] Pearson, P.N. 1992 Survivorship Analysis of Fossil Taxa when Real-Time Extinction Rates Vary: The Paleogene Planktonic Foraminifera. *Paleobiology* **18**, 115-131. (doi:10.2307/2400994).
- [60] Simpson, G.G. 1951 The Species Concept. *Evolution* **5**, 285-298. (doi:10.2307/2405675).
- [61] Wiley, E.O. 1978 The Evolutionary Species Concept Reconsidered. *Syst. Biol.* **27**, 17-26. (doi:10.2307/2412809).
- [62] Ezard, T.H.G., Pearson, P.N., Aze, T. & Purvis, A. 2012 The meaning of birth and death (in macroevolutionary birth-death models). *Biol. Letters* **8**, 139-142. (doi:10.1098/rsbl.2011.0699).
- [63] Chaudhary, C., Saeedi, H. & Costello, M. In review Bimodality of latitudinal gradients in marine species richness. *Trends in Ecology & Evolution*.
- [64] Mittelbach, G.G., Schemske, D.W., Cornell, H.V., Allen, A.P., Brown, J.M., Bush, M.B., Harrison, S.P., Hurlbert, A.H., Knowlton, N., Lessios, H.A., et al. 2007 Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecol. Lett.* **10**, 315-331. (doi:10.1111/j.1461-0248.2007.01020.x).
- [65] Roy, K. & Witman, J.D. 2009 Spatial patterns of species diversity in the shallow marine invertebrates: Patterns, processes, and prospects. In *Marine Macroecology* (eds. J.D. Witman & K. Roy), pp. 101-121, The University of Chicago Press.
- [66] Briggs, J.C. 2006 Proximate sources of marine biodiversity. *J Biogeogr* **33**, 1-10. (doi:10.1111/j.1365-2699.2005.01374.x).
- [67] Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. 2004 Toward a metabolic theory of ecology. *Ecology* **85**, 1771-1789. (doi:10.1890/03-9000).
- [68] Colwell, R.K. & Hurtt, G.C. 1994 Nonbiological gradients in species richness and a spurious Rapoport effect. *Am. Nat.* **144**, 570-595. (doi:10.2307/2462939).
- [69] Yasuhara, z., Hunt, G., Dowsett, H.J., Robinson, M.M. & Stoll, D.K. 2012 Latitudinal species diversity gradient of marine zooplankton for the last three million years. *Ecol. Lett.* **15**, 1174-1179. (doi:10.1111/j.1461-0248.2012.01828.x).
- [70] Lamanna, C., Blonder, B., Violle, C., Kraft, N.J.B., Sandel, B., Šímová, I., Donoghue, J.C., Svenning, J.-C., McGill, B.J., Boyle, B., et al. 2014 Functional trait space and the latitudinal diversity gradient. *Proc. Natn. Acad. Sci.* **111**, 13745-13750. (doi:10.1073/pnas.1317722111).
- [71] Villalobos, F., Carotenuto, F., Raia, P. & Diniz-Filho, J.A.F. this volume Phylogenetic fields through time: temporal dynamics of geographic co-occurrence and phylogenetic structure within species ranges. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*.
- [72] Morey, A.E., Mix, A.C. & Pisias, N.G. 2005 Planktonic foraminiferal assemblages preserved in surface sediments correspond to multiple environment variables. *Quaternary Sci. Rev.* **24**, 925-950.
- [73] Darling, K.F., Wade, C.M., Stewart, I.A., Kroon, D., Dingle, R. & Brown, A.J.L. 2000 Molecular evidence for genetic mixing of Arctic and Antarctic subpolar populations of planktonic foraminifers. *Nature* **405**, 43-47.
- [74] Zachos, J.C., Dickens, G.R. & Zeebe, R.E. 2008 An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature* **451**, 279-283.
- [75] Cervato, C. & Burckle, L. 2003 Pattern of first and last appearance in diatoms: Oceanic circulation and the position of polar fronts during the Cenozoic. *Paleoceanography* **18**, 1055. (doi:10.1029/2002PA000805).
- [76] Pagani, M., Huber, M., Liu, Z., Bohaty, S.M., Henderiks, J., Sijp, W., Krishnan, S. & de Conto, R.M. 2011 The role of carbon dioxide during the onset of Antarctic glaciation. *Science* **334**, 1261-1264. (doi:10.1126/science.1203909).

- [77] John, E.H., Pearson, P.N., Coxall, H.K., Birch, H., Wade, B.S. & Foster, G.L. 2013 Warm ocean processes and carbon cycling in the Eocene. *Phil. Trans. R. Soc. A* **371**. (doi:10.1098/rsta.2013.0099).
- [78] Pearson, P.N. & Coxall, H.K. 2014 Origin of the Eocene planktonic foraminifer *Hantkenina* by gradual evolution. *Palaeontology* **57**, 243-267. (doi:10.1111/pala.12064).
- [79] Boersma, A. & Premoli Silva, I. 1991 Distribution of Paleogene planktonic foraminifera — analogies with the Recent? *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **83**, 29-47.
- [80] Pearson, P.N., Olsson, R.K., Huber, B.T., Hemleben, C. & Berggren, W.A. 2006 *Atlas of Eocene Planktonic Foraminifera*, Cushman Foundation; 514 p.
- [81] Coxall, H.K., Wilson, P.A., Pearson, P.N. & Sexton, P.F. 2007 Iterative evolution of digitate planktonic foraminifera. *Paleobiology* **33**, 495-516. (doi:10.1666/06034.1).
- [82] Coxall, H.K., Pearson, P.N., Shackleton, N.J. & Hall, M.A. 2000 *Hantkeninid* depth adaptation: An evolving life strategy in a changing ocean. *Geology* **28**, 87-90. (doi:10.1130/0091-7613(2000)28<87:hdaael>2.0.co;2).
- [83] Pearson, P.N., McMillan, I.K., Wade, B.S., Dunkley Jones, T., Coxall, H.K., Bown, P.R. & Lear, C.H. 2008 Extinction and environmental change across the Eocene-Oligocene boundary in Tanzania. *Geology* **36**, 179-182. (doi:10.1130/G24308a.1).
- [84] Lazarus, D. 1994 Neptune: A marine micropaleontology database. *Math Geol* **26**, 817-832. (doi:10.1007/BF02083119).
- [85] Spencer-Cervato, C. 1999 The Cenozoic Deep Sea Microfossil Record: Explorations of the DSDP/ODP Sample Set Using the Neptune Database. *Palaeontol. Electron.* **2**, 268pp.
- [86] Inglis, G.N., Farnsworth, A., Lunt, D., Foster, G.L., Hollis, C.J., Pagani, M., Jardine, P.E., Pearson, P.N., Markwick, P., Galsworthy, A.M.J., et al. 2015 Descent toward the Icehouse: Eocene sea surface cooling inferred from GDGT distributions. *Paleoceanography*, n/a-n/a. (doi:10.1002/2014PA002723).
- [87] Berger, W. 1970 Planktonic foraminifera: Selective solution and the lysocline. *Mar. Geol.* **8**, 111-138. (doi:10.1016/0025-3227(70)90001-0).
- [88] Wood, S.N. 2006 *Generalized Additive Models: An introduction with R*, Chapman & Hall/CRC; 392 p.
- [89] Lunt, D.J., Dunkley Jones, T., Heinemann, M., Huber, M., LeGrande, A., Winguth, A., Loptson, C., Marotzke, J., Roberts, C.D., Tindall, J., et al. 2012 A model–data comparison for a multi-model ensemble of early Eocene atmosphere–ocean simulations: EoMIP. *Clim. Past* **8**, 1717-1736. (doi:10.5194/cp-8-1717-2012).
- [90] Dunkley Jones, T., Lunt, D.J., Schmidt, D.N., Ridgwell, A., Sluijs, A., Valdes, P.J. & Maslin, M. 2013 Climate model and proxy data constraints on ocean warming across the Paleocene–Eocene Thermal Maximum. *Earth Sci. Rev.* **125**, 123-145.
- [91] Lunt, D.J., Valdes, P.J., Jones, T.D., Ridgwell, A., Haywood, A.M., Schmidt, D.N., Marsh, R. & Maslin, M. 2010 CO₂-driven ocean circulation changes as an amplifier of Paleocene–Eocene thermal maximum hydrate destabilization. *Geology* **38**, 875-878. (doi:10.1130/g31184.1).
- [92] Lunt, D.J., Farnsworth, A., Loptson, C., Foster, G.L., Markwick, P., O'Brien, C.L., Pancost, R.D., Robinson, S.A. & Wrobel, N. 2015 Palaeogeographic controls on climate and proxy interpretation. *Clim. Past Discuss.* **11**, 5683-5725. (doi:10.5194/cpd-11-5683-2015).
- [93] R Core Team. 2014 R: A language and environment for statistical computing. (R Foundation for Statistical Computing.
- [94] Crawley, M.J. 2007 *The R book*, John Wiley & Sons; 942 p.
- [95] Burnham, K.P. & Anderson, D.R. 2002 *Model Selection and Multimodel Inference: A Practical Information-Theoretical Approach*, Springer, New York.
- [96] Pearson, P.N., van Dongen, B.E., Nicholas, C.J., Pancost, R.D., Schouten, S., Singano, J.M. & Wade, B.S. 2007 Stable warm tropical climate through the Eocene Epoch. *Geology* **35**, 211-214. (doi:10.1130/g23175a.1).

- [97] Bijl, P.K., Schouten, S., Sluijs, A., Reichart, G.-J., Zachos, J.C. & Brinkhuis, H. 2009 Early Palaeogene temperature evolution of the southwest Pacific Ocean. *Nature* **461**, 776-779.
- [98] Huber, M. & Caballero, R. 2011 The early Eocene equable climate problem revisited. *Clim. Past* **7**, 603-633. (doi:10.5194/cp-7-603-2011).
- [99] Sagoo, N., Valdes, P., Flecker, R. & Gregoire, L.J. 2013 The Early Eocene equable climate problem: can perturbations of climate model parameters identify possible solutions? *Phil. Trans. R. Soc. A* **371**.
- [100] Kiehl, J.T. & Shields, C.A. 2013 Sensitivity of the Palaeocene–Eocene Thermal Maximum climate to cloud properties. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* **371**.
- [101] John, E.H., Wilson, J.D., Pearson, P.N. & Ridgwell, A. 2014 Temperature-dependent remineralization and carbon cycling in the warm Eocene oceans. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **413**, 158-166.
- [102] Olivarez Lyle, A. & Lyle, M.W. 2006 Missing organic carbon in Eocene marine sediments: Is metabolism the biological feedback that maintains end-member climates? *Paleoceanography* **21**, 1-13. (doi:10.1029/2005pa001230).
- [103] Peters, S.E. 2008 Environmental determinants of extinction selectivity in the fossil record. *Nature* **454**, 626-629.
- [104] Magurran, A.E. 2004 *Measuring biological diversity*, Blackwell Publishing Ltd; 256 p.
- [105] Purvis, A. & Hector, A. 2000 Getting the measure of biodiversity. *Nature* **405**, 212-219.
- [106] Mouchet, M.A., Villéger, S., Mason, N.W.H. & Moullot, D. 2010 Functional diversity measures: an overview of their redundancy and their ability to discriminate community assembly rules. *Funct. Ecol.* **24**, 867-876. (doi:10.1111/j.1365-2435.2010.01695.x).
- [107] Monnet, A.-C., Jiguet, F., Meynard, C.N., Moullot, D., Mouquet, N., Thuiller, W. & Devictor, V. 2014 Asynchrony of taxonomic, functional and phylogenetic diversity in birds. *Global Ecol. Biogeogr.* **23**, 780-788. (doi:10.1111/geb.12179).
- [108] de Carvalho, R.A. & Tejerina-Garro, F.L. 2015 Relationships between taxonomic and functional components of diversity: implications for conservation of tropical freshwater fishes. *Freshwater Biol.* **60**, 1854-1862. (doi:10.1111/fwb.12616).
- [109] Cadotte, M., Albert, C.H. & Walker, S.C. 2013 The ecology of differences: assessing community assembly with trait and evolutionary distances. *Ecol. Lett.* **16**, 1234-1244. (doi:10.1111/ele.12161).
- [110] Li, G. & Chen, Z. 1985 Projection-Pursuit Approach to Robust Dispersion Matrices and Principal Components: Primary Theory and Monte Carlo. *J Am Stat Assoc* **80**, 759-766. (doi:10.1080/01621459.1985.10478181).
- [111] Ezard, T., Pearson, P. & Purvis, A. 2010 Algorithmic approaches to aid species' delimitation in multidimensional morphospace. *BMC Evo. Biol.* **10**, 175.
- [112] Singhal, A. & Seborg, D.E. 2005 Clustering multivariate time-series data. *J Chemometr* **19**, 427-438. (doi:10.1002/cem.945).
- [113] Berggren, W.A. 1969 Rates of Evolution in Some Cenozoic Planktonic Foraminifera. *Micropaleontology* **15**, 351-365. (doi:10.2307/1484931).